

Pistachio

sonar



soil



foliar

Special for pistachio

Composition

%w/w

Magnesium (MgO)	2,0
Copper (Cu)	2,0
Zinc (Zn)	1,0
Sulfur (SO ₃)	7,4



Characteristics

Micronutrients complexed with Gluconic acid in liquid formulation indicated to prevent deficiencies in pistachio crops.

Pistachio Sonar has been developed for preventive and curative application on foliar spray or fertirrigation. Its use is suggested as supplement to a regular, balanced fertilizer program to enhance yields and improve quality applications.

Foliar application of **Pistachio Sonar** may be particularly beneficial during periods of peak nutrient demand, for crops grown on soils having poor nutrient availability or to crops suffering from a weakened root system.



YIELD AND QUALITY

Application



FOLIAR APPLICATION	DOSE
Foliar	200-400 cc/100L H ₂ O
Separate applications	7-14 days.



SOIL APPLICATION	DOSE
Fertility Maintenance	2-5 L/ha
Mild deficiencies	4-9 L/ha
Serious deficiencies	8-20 L/ha

Actively growing plants with young leaves absorb foliarly applied most efficiently. Immediate post-irrigation or post-rainfall application will enhance foliar uptake. Thorough coverage and wetting are needed for optimum foliar spray results. Application should be separated at minimum of 7-14 days. Avoid Foliar applications when plants are under moisture stress. AVOID foliar application to deciduous tree crops during bloom. Use up to 2 L of in a minimum of 200 L of water per hectare.

Shake, stir, or swirl contents before using. Always add to the spray tank before adding pesticides. Pistachio Sonar is compatible with insecticides and acaricides normally under use. In case of doubt, a previous test is advisable. Pistachio Sonar is not recommended to be mixed with highly alkaline products. Soil applications are recommended with nitrogen fertilizer solutions.



SHAKE WELL BEFORE USE



IMPORTED FROM EU

