

Composition

	%w/v
Gibberellic acid (GA ₃)	1,6
Soluble liquid (SL)	



System and time of application

Ensure that the wind does not drag the spraying to other neighboring crops. Apply the product shortly after mixing with water to prevent decomposition.

In the event of a precipitation, at 8 hours after treatment, it will lose some of its effectiveness, it is advisable to repeat the treatment.

Usage instructions

Treat with high water volume (600-1500 L / Ha). Add wetting

Characteristics

sonar GIB is a plant growth regulator characterized by its physiological and morphological effects. Acts at very low concentrations; is translocated inside the plant and usually affects only the aerial parts.

sonar GIB reinforces apical dominance, stimulating flowering, fruiting set, breaking the dormancy of seeds and vegetative organs and removing stress from some virus.

Growth, blooming and fruiting

Application

Crop	Doses (cc/hl)	Treatment
Artichoke	40	To induce growth and harvest earliness. Treat to start fruiting
Clementine	30 - 50	To improve the consistency of the peel (when the green treat is gone)
Lemon and clementine boneless	40	To induce fruit set and fruit setting. Treat at petal drop and repeat the treatment at 3-4 weeks
Pear (cv. Blanquilla)	60 - 70	To reduce fruit drop during filling and prevent frost damage. Treat with 30-60% open flower or 48 hours to avoid the effect of frost
Tangerine	40	To prevent fruit drop (treat at petal drop, repeating at 3-4 weeks) and improve the consistency of the peel (when the green treat is gone)
Vine (cv. Macabeo)	30 - 50	To induce the elongation of the cluster and peduncles of the fruit. Apply before the falling of the flowerhoods
Strawberry	60	To promote fruit set and fruit growth. Treat a little before the start of flowering. Can be reduced the dose in plantations over a year old

