

SONAI Gluco-Ca

COMPLEXED ORGANIC CALCIUM CORRECTOR





COMPOSITION	%w/w
Calcium (CaO) Water soluble	8,0
Density: 1.2	
pH: 9-10	



CHARACTERISTICS

SONAR GLUCO Ca is a gluco-complexed liquid fertilizer for use as a foliar feed to maintain or increase calcium levels in plants.

SONAR GLUCO Ca is specifically designed to provide Calcium to fruit and vegetable crops more efficiently than other forms of Calcium. Gluconic acid complexes calcium ion enabling it to move into the plant via the phloem.

SONAR GLUCO Ca complex reaches the fruit forming tissue, the sugar bond breaks down and the Calcium flows to where it is needed.

Unlike Calcium Chloride and Calcium Nitrate, SONAR GLUCO Ca will not produce injuries to the foliage and fruit, such as burned leaves and spotted fruit enabling **SONAR GLUCO Ca** to be used during the growing season.

ADVANTAGES

Increases in number of leaves



Better/increase dry

DOSES AND APPLICATION

Crop	Aim / problem	Recommendation	Time
Cereals	Vitality, stalk stability	1-3 times 5 I/ha	From the beginning of tillering.
Citrus fruits	Vitality, fruit firmness, storage and transport stability.	2-5 times 5 I/ha	From fruit set.
General Vegetables	Vitality, fruit strength, storage and transport stability, against internal fire, margin necrosis and flower rot.	2-5 times 5-10 l/ha	Once sufficient leaf mas had developed or from fruit set to harvest.
In all crops	For calcium supply, cell wall strength, reduction of radiation stress (anntioxidant), improvement of fruit quality and storage stability	5-10 I/ha (for leaf fertilisation with at least 500 litres of water. In case of application with the backpack sprayer 1%. Only in chloride-insensitive Cultures and not during flowering!)	When required
Oilseed rape	Vitality, stalk stability	1-3 times 5- 10 I/ha	From 4-leaf stage
Ornamental plants	Vitality, leaf quality, transport stability.	1-3 times 5 I/ha.	Once sufficient leaf mas has developed.
Pome fruit	Vitality, fruit firmness, storage and transport stability.	4-6 times 5-10 I/ha.	From walnut size to harvesting.
Potatoes	Tuber and skin quality, improvement in storage life.	2-4 times 5 I/ha	From beginning of row closure.
Stone fruit	Vitality, fruit firmness, storage and transport stability.	2-5 times 5-10 I/ha.	From fruit set.
Strawberries	Vitality, fruit firmness, storage and transport stability.	2-4 times 5 I/ha.	From fruit set
Sugar beet	Quality, storage and transport stability.	1-3 times 5 I/ha	From 6- leaf stage.
Sunflowers	Vitality, stalk stability	1-3 times 5 I/ha	From 4-leaf stage
Table grapes	Vitality, berry skin firmness, storage and transport stability.	2-5 times 5 I/ha	Pea size to harvesting.
Wine grapes	Vitality, berry skin firmness, storage and transport stability	2-5 times 5 I/ha	Pea size to harvesting.

Shake it before use

CAUTION: check compatibility with standard jar test.





