COMPLEX DENSO

NPK Fertilizer with trace elements. Gel formulation





Composition	%w/v
Total nitrogen (N)	10,0
Phosphorus (P ₂ O ₅)	50,0
Potassium (K,Ô)	10,0
Boron (B)	0.016
Iron (Fe) chelating agent EDTA	0,047
Copper (Cu) chelating agent EDTA	0,016
Manganese (Mn) chelating agent EDTA	0,016
Zinc (Zn) chelating agent EDTA	0,016
Density 1 35	





GREEN GEL

10-50-10 +Te

Neutral pH , unlike most liquid foliar that are highly acidic or highly alkaline. COMPLEX DENSO can be used at higher doses, not being aggressive with the cells that form stomas

More comfortable for the farmer to dosing per volume instead of on weight

Best solution in terms of speed and ease to use. Allow higher liquid dispersion homogeneity than solid products

Guarantee solubility by its

Adjuvant: promotes effectiveness of plant protection products when applied jointly

nutritional product and not just a simple mixture of raw materials, as are most of NPK fertilizers in powder form.

and simultaneous solubility of all the nutrients, during use, while avoiding sedimentation in the storage containers of the nutrient solution. In contrast, common NPK water soluble powder fertilizers, which are produced through a mixture of raw materials, have increase variability in grain size that results in a non-uniform dilution of nutrients, since the smallest grains are dissolved firstly.

The conductivity and the salinity index are maintained in very low levels so that the soil will not be burndened with undesirable, salt concentration.

Application is suitable for different crops: fruit trees, coffee, olive trees, vegetable crops, industrial crops, meadows, etc. It can be used in drip irrigation, foliar application and food irrigation.

recommend a trial or consult our technical department.

Crops	Dosages	Applications
Cereals	2-5l/ha 250 ml/100l	Early in crop cyrcle. Followed by 2nd application 14 days later.
Paprika	2,0 - 3,0l/ha 200-300 ml/100l	1st application 3 weeks after transplanting, followed with a 2nd application 14 days later.
Roses and Ornamentals	2,0 - 3,0l/ha 200-300 ml/100l	Monthly applications on perennials. 2 applications 14 days apart on annual during initial growth stages.
Strawberries	3,0l/ha 300 ml/100l	Single application 3 weeks after planting.
Tomatos and Peppers	2,0 - 3,0l/ha 200-300 ml/100l	1st application 3 weeks after transplanting, followed by a 2nd application 14 days later.
Vegetables	3,0l/ha 200-300 ml/100l	1 to 2 applications early on in growth period of crop.
Other crops	2,0 - 2,5 l/ha 300 ml/100l	For crops with phosphate deficiencies, repeat at 10-14 days intervals as required.



FFRTII IZFR









It is recommended to shake before use.

