

Composition	%w/w
Iron (Fe)	7,50
Manganese (Mn)	3,50
Zinc (Zn)	0,70
Boron (B)	0,65
Copper (Cu)	0,28
Molybdenum (Mo)	0,03
Chelating Agent: EDTA	



Characteristics

kelat MIX micro is a solid compound, highly-soluble in all types of water and whose Iron, Manganese, Copper, Zinc, Boron, Molybdenum and Magnesium micronutrients contribute simultaneously to the plant by providing the necessary dosage of nutrients that are indispensable for the perfect development of any crop.

kelat MIX micro allows easy and correct dosage which offers instant solubility and high agronomic efficiency; obtaining good yields and high quality crops.

Except for the Boron and Molybdenum, the nutrients in **kelat MIX micro** are included in a molecule (EDTA, ethylenediaminetetraacetic acid) that protects them in the soil and, when applied to the leaves, facilitates their uptake and transport to the plant.

Its unique manufacturing process obtained by chemical mixing in the liquid phase ensures a complete chelation and a total homogeneity; keeping the same composition, size, density, color and guaranteed nutritional balance in each microgranule.

Application

Crops	Dosage	Treatment
Fertirrigation		
General dose	3 - 4 Kg/Ha	Every 7-10 days during the crop cycle
Hydroponic		
General dose	0,3 - 0,5 Kg/m ³ of water	Prepare a stock solution 100 times concentrated and employ 1L each 100 L of water irrigation. Use the larger doses during periods of increased growth of crops.
Foliar		
General dose	1 - 1,5 Kg/Ha or 100 - 150 g/100L	Applied when symptoms appear.
Horticultural	3x75 - 100 g/hl of water (3x0,5 - 1 Kg/Ha)	At 10-15 day intervals, beginning when the foliage is enough.
Fruit trees, vines, citrus and olive trees	100 g/100 L of water (1 Kg/Ha) 100 g/100 L of water (1 - 1,5Kg/Ha)	First bloom After fruit set
Field crops, industrial crops	1 Kg/Ha	During the crop cycle
Potatoes and vegetables Bulb	4 x 1 Kg/Ha	At 7/10 days intervals, starting at 10 cm of growth. Apply in a minimum of 500 L / Ha water.
Ornamental	75 - 150 g/hl of water (0,5 - 1,5 Kg/Ha)	2-4 applications with intervals of 7-10 days at the beginning of the growing season

FERTILIZER



IMPORTED FROM EU

