

# Sonar ilicFulvic

Silicon and Calcium  
fertilizer with Fulvic acids



Composition	%w/w
Silicon (SiO <sub>2</sub> )	7,0
Calcium Oxide (CaO)	7,0
Fulvic acids	14,5

## Characteristics

SONAR SILIC FULVIC is a product designed to provide crops Ca and Si. It also incorporates Fulvic Acids that act as effective synergists in the uptake and transport of Ca and Si within the plant.

### Improved cell structure strength

reducing the incidence of lodging. Si accumulates in the epidermal tissues increasing the mechanical stability of the plant.

### Improved resistance to pathogens and insects

Si deposition in the epidermis tissues provides a physical barrier to pathogens and insects.

### Improved photosynthetic activity.

The improved structure of the plant has been shown to improve its ability to capture the light.

### Reduced drought and heat stress

The deposition of Si in the plant tissues reduces transpiration rates.

**Reduced salt stress** by inhibiting sodium absorption.

**Improved utilization of applied fertilizers** particularly Nitrogen, Phosphorus and Potassium.

**Alleviates toxicity** of Iron, Manganese, Cadmium and Aluminium

NUTRITION

ENHANCE PLANT GROWTH

INCREASES RESISTANCE  
TO DISEASES

QUALITY

## Applications

Crops	Rate/ha 300L water	Details
Beans, Carrots, Celery, Lettuce, Brassicas	2-3L/ha	Apply at 2 leaf stage. Apply 2-4 days after sowing via solid set Apply as a soil drench at transplant or emergence. Repeat 7-10 days later.
Citrus	5-7L/ha	Apply to juvenile trees at early establishment - repeat as necessary. Mature trees-treat at spring and autumn growth flush.
Cucurbits, Cut flower production and Bulb production	2-3L/ha	Apply at 1 - 2 leaf stage - repeat application at 2 - 4 leaf stage. Apply at emergence or transplant. Drench bulb at planting. Repeat 2 weeks after emergence. Continue if weak stem symptoms are evident.

Crops	Rate/ha 300L water	Details
General production	2-3L/ha	Apply at emergence or transplant - repeat 7 - 10 day intervals as required.
Trickle irrigation	5-7L/ha	1:300 minimum for trickle.
Top fruit Potatoes	2-3L/ha	Apply at transplant - repeat as required during establishment. Apply 1 week after planting - repeat at 7 - 10 day intervals.
Maize, Cereals and other field crops	3-5L/ha	Apply when leaf area is sufficient to intercept foliar spray. Silica treatments can reduce droopy growth and lodging.
Tomatoes/ Capsicum	2-3L/ha	Apply at transplanting - trickle or foliar. Mature plants: repeat when stalk weakness is evident.
Vines		Apply at vine establishment, repeat at flower truss visible.

## Silicon and postharvest life or produce

Researchers have shown that Silicon can inhibit ethylene which reduces the speed of aging and death of harvested plant parts. Silicon treated plant have also been shown to maintain their chlorophyll (green) content over a longer period. The end result is produce with better shelf life and appearance.



IMPORTED  
FROM EU



sonar  
agro