


Composition

	%w/w
Calcium (CaO)	55,00
Magnesium (MgO)	0,15



QUALITY AND HEALTH IN PRE-HARVEST



SUNBURN EFFECT



APPLIED PRODUCT

Application

Crops	Doses	Remarks
FRUIT TREES: Apple trees, Pear trees, Lemon, Orange, Tangerine, Clementine, Grapefruit, Olives, Peaches, Nectarines, Pomegranates, Persimmons, Avocado	5-10 Kg/100 L water	Apply in aqueous solutions in a traditional way, with nebulizer. It is recommended to apply on two consecutive passes and in opposite directions. It is necessary that the tree is completely covered (homogeneous distribution) and white color. Make 3-5 applications every 7 days maximum. These applications should be initiated before the period of maximum susceptibility. Use wetting from 1500 to 3000 L / ha
VEGETABLES: Tomatoes, Peppers, Melon, Watermelon	4-7 Kg/100 L water	It's recommended to apply on a volume of 600L/ha two consecutive passes in opposite directions. Apply during periods of higher susceptibility corresponding to the start of veraison when the fruit begins to change from green to orange.

Application time: applications should begin when temperatures exceed the thermal threshold established by the technicians of the area.
 Frequency of application: every 20 to 30 days, depending on weather conditions and/or rate of growth of the fruit.
 Number of applications: 3-4 applications per season and depending on weather conditions.

1 Kg 5 Kg 20 Kg

